

# CONTRACTING QUADRATIC OPERATORS OF BISEXUAL POPULATION

Nasir N. Ganikhodjaev

Department of Computational and Theoretical Sciences, Faculty of Science, IIUM,  
25200 Kuantan, Malaysia.

Email: nasirgani@hotmail.com

Uygun U. Jamilov

Department of Computational and Theoretical Sciences, Faculty of Science, IIUM,  
25200 Kuantan, Malaysia.

Institute of Mathematics at the National University Uzbekistan, Tashkent,  
Uzbekistan,

e-mail: jamilovu@yandex.ru, ujamilov@rambler.ru

**Abstract.** In this paper we find a sufficient condition under which the operator of bisexual population is contraction and show that this condition is not necessary.

**Mathematics Subject Classification(2000):** Primary 37N25, Secondary 92D10.

**Key words.** Quadratic stochastic operator, fixed point, trajectory, contracting operators.

## 1 Introduction

The action of genes is manifested statistically in sufficiently large communities of matching individuals (belonging to the same species). These communities are called populations [2]. The population exists not only in space but also in time, i.e. it has its own life cycle. The basis for this phenomenon is reproduction by mating. Mating in a population can be free or subject to certain restrictions.

The whole population in space and time comprises discrete generations  $F_0, F_1, \dots$ . The generation  $F_{n+1}$  is the set of individuals whose parents belong to the  $F_n$  generation. A state of a population is a distribution of probabilities of the different types of organisms in every generation. Type partition is called differentiation. The simplest example is sex differentiation. In bisexual population any kind of differentiation must agree with the sex differentiation, i.e. all the organisms of one type must belong to the same sex. Thus, it is possible to speak of male and female types.

The evolution (or dynamics) of a population comprises a determined change of state in the next generations as a result of reproductions and selection. This evolution of a population can be studied by a dynamical system (iterations) of a quadratic stochastic operator.

The history of the quadratic stochastic operators can be traced back to the work of S. Bernshtein [1]. For more than 80 years this theory has been developed and many papers were published (see [1]-[15]). Several problems of physical and biological systems lead to necessity of study the asymptotic behavior of the trajectories of quadratic stochastic operators.

Let  $E = \{1, 2, \dots, m\}$ . By the  $(m-1)$ -simplex we mean the set

$$S^{m-1} = \{\mathbf{x} = (x_1, \dots, x_m) \in R^m : x_i \geq 0, \sum_{i=1}^m x_i = 1\}. \quad (1)$$

Each element  $\mathbf{x} \in S^{m-1}$  is a probability measure on  $E$  and so it may be looked upon as the state of a biological (physical and so on) system of  $m$  elements.

A quadratic stochastic operator  $V : S^{m-1} \rightarrow S^{m-1}$  has the form

$$V : x'_k = \sum_{i,j=1}^m p_{ij,k} x_i x_j, \quad (k = 1, \dots, m), \quad (2)$$

where  $p_{ij,k}$ —coefficient of heredity and

$$p_{ij,k} = p_{ji,k} \geq 0, \quad \sum_{k=1}^m p_{ij,k} = 1, \quad (i, j, k = 1, \dots, m). \quad (3)$$

For a given  $x^{(0)} \in S^{m-1}$ , the trajectory  $\{x^{(n)}\}$ ,  $n = 0, 1, 2, \dots$  of  $x^{(0)}$  under the action of QSO (2) is defined by  $x^{(n+1)} = V(x^{(n)})$ , where  $n = 0, 1, 2, \dots$

One of the main problems in mathematical biology is to study the asymptotic behavior of the trajectories. There are many papers devoted to study of the evolution of the free population, i.e. to study of dynamical system generated by quadratic stochastic operator (2), see e.g. [3]-[14]. In [13] a survey of theory quadratic stochastic operators is given.

In this paper we find a condition under which the evolutionary operators of bisexual population is contraction.

## 2 Definitions

In this section following [2], we describe the evolution operator of a bisexual population. Assuming that the population is bisexual we suppose that the set of females can be partitioned into finitely many different types indexed by  $\{1, 2, \dots, n\}$  and, similarly, that the male types are indexed by  $\{1, 2, \dots, \nu\}$ . The number  $n + \nu$  is called the dimension of the population. The population is described by its state vector  $(\mathbf{x}, \mathbf{y})$  in  $S^{n-1} \times S^{\nu-1}$ , the product of two unit simplexes in  $\mathbb{R}^n$  and  $\mathbb{R}^\nu$  respectively.

Vectors  $\mathbf{x}$  and  $\mathbf{y}$  are the probability distributions of the females and males over the possible types:

$$x_i \geq 0, \sum_{i=1}^n x_i = 1; \quad y_j \geq 0, \sum_{j=1}^{\nu} y_j = 1. \quad (4)$$

Denote  $S = S^{n-1} \times S^{\nu-1}$ . We call the partition into types hereditary if for each possible state  $\mathbf{z} = (\mathbf{x}, \mathbf{y}) \in S$  describing the current generation, the state  $\mathbf{z}' = (\mathbf{x}', \mathbf{y}') \in S$  is uniquely defined describing the next generation. This means that the association  $\mathbf{z} \rightarrow \mathbf{z}'$  defined a map  $V : S \rightarrow S$  called the evolution operator. For any point  $\mathbf{z}^{(0)} \in S$  the sequence  $\mathbf{z}^{(t)} = V(\mathbf{z}^{(t-1)}), t = 1, 2, \dots$  is called the trajectory of  $\mathbf{z}^{(0)}$ . Let  $p_{ij,k}^{(f)}$  and  $p_{ij,l}^{(m)}$  be inheritance coefficients defined as the probability that a female offspring is type  $k$  and, respectively, that a male offspring is type  $l$ , when the parental pair is  $ij (i, k = 1, 2, \dots, n; \text{ and } j, l = 1, 2, \dots, \nu)$ . We have

$$p_{ij,k}^{(f)} \geq 0, \sum_{k=1}^n p_{ij,k}^{(f)} = 1, \quad p_{ij,l}^{(m)} \geq 0, \sum_{l=1}^{\nu} p_{ij,l}^{(m)} = 1. \quad (5)$$

Let  $\mathbf{z}' = (\mathbf{x}', \mathbf{y}')$  be the state of the offspring population at the birth stage. This is obtained from inheritance coefficients as

$$W : \begin{cases} x'_k = \sum_{i,j=1}^{n,\nu} p_{ij,k}^{(f)} x_i y_j, & (1 \leq k \leq n) \\ y'_l = \sum_{i,j=1}^{n,\nu} p_{ij,l}^{(m)} x_i y_j, & (1 \leq l \leq \nu). \end{cases} \quad (6)$$

We see from (6) that for a bisexual population the evolution operator is a quadratic mapping of  $S$  into itself. But for free population the operator is quadratic mapping of the simplex into itself given by (2).

In [8] an algebra of the bisexual population is defined as the following: Consider  $\{e_1, \dots, e_{n+\nu}\}$  the canonical basis on  $\mathbb{R}^{n+\nu}$  and divide the basis as  $e_i^{(f)} = e_i, i = 1, \dots, n$  and  $e_i^{(m)} = e_{n+i}, i = 1, \dots, \nu$ . Introduce on  $\mathbb{R}^{n+\nu}$  a multiplication defined by

$$\begin{aligned} e_i^{(f)} e_j^{(m)} &= e_j^{(m)} e_i^{(f)} = \frac{1}{2} \left( \sum_{k=1}^n p_{ij,k}^{(f)} e_k^{(f)} + \sum_{l=1}^{\nu} p_{ij,l}^{(m)} e_l^{(m)} \right); \\ e_i^{(f)} e_k^{(f)} &= 0, \quad i, k = 1, \dots, n; \\ e_j^{(m)} e_l^{(m)} &= 0, \quad j, l = 1, \dots, \nu; \end{aligned} \quad (7)$$

Thus the coefficients of bisexual inheritance is the structure constants of an algebra, i.e. a bilinear mapping of  $\mathbb{R}^{n+\nu} \times \mathbb{R}^{n+\nu}$  to  $\mathbb{R}^{n+\nu}$ . The general formula for the multiplication is the extension of (7) by bilinearity, i.e. for  $z, t \in \mathbb{R}^{n+\nu}$ ,

$$z = (x, y) = \sum_{i=1}^n x_i e_i^{(f)} + \sum_{j=1}^{\nu} y_j e_j^{(m)}, \quad t = (u, v) = \sum_{i=1}^n u_i e_i^{(f)} + \sum_{j=1}^{\nu} v_j e_j^{(m)}$$

using (7), we obtain

$$zt = \frac{1}{2} \sum_{k=1}^n \left( \sum_{i=1}^n \sum_{j=1}^{\nu} p_{ij,k}^{(f)} (x_i v_j + u_i y_j) \right) e_k^{(f)} + \frac{1}{2} \sum_{l=1}^{\nu} \left( \sum_{i=1}^n \sum_{j=1}^{\nu} p_{ij,l}^{(m)} (x_i v_j + u_i y_j) \right) e_l^{(m)}. \quad (8)$$

From (8) and using (6), in the particular case that  $z = t$ , i.e.  $x = u$  and  $y = v$ , we obtain

$$zz = z^2 = \sum_{k=1}^n \left( \sum_{i=1}^n \sum_{j=1}^{\nu} p_{ij,k}^{(f)} x_i y_j \right) e_k^{(f)} + \sum_{l=1}^{\nu} \left( \sum_{i=1}^n \sum_{j=1}^{\nu} p_{ij,l}^{(m)} x_i y_j \right) e_l^{(m)} = W(z).$$

for any  $z \in S$ . This algebraic interpretation is very useful. For example, a bisexual population state  $z = (x, y)$  is an equilibrium (fixed point) precisely when  $z$  is an idempotent element of the set  $S$ , i.e.  $z = z^2$ .

The algebra  $\mathcal{B} = \mathcal{B}_W$  generated by the evolution operator  $W$  (see (6)) is called the *evolution algebra of the bisexual population*.

In [8] it was shown that if  $z$  is a fixed point then  $z \in R_0^{n+\nu} \cup R_1^{n+\nu}$ , where

$$R_\eta^{n+\nu} = \{z = (x, y) : \sum_{i=1}^n x_i = \sum_{j=1}^{\nu} y_j = \eta\}, \quad \eta = 0, 1. \quad (9)$$

For simplex  $S = S^{n-1} \times S^{\nu-1}$  by tangent space we get

$$R_0^{n+\nu} = \{z = (x, y) : \sum_{i=1}^n x_i = \sum_{j=1}^{\nu} y_j = 0\}. \quad (10)$$

### 3 Contracting operators

In operator theory, a bounded operator  $W : X \rightarrow Y$  between normed vector spaces  $X$  and  $Y$  is said to be a *contraction* if its operator norm  $\|W\| \leq 1$ .

An extremal example of a quadratic contraction is the constant operator. In this case the coefficients  $p_{ij,k}^{(f)}, p_{ij,l}^{(m)}$  do not depend on  $i$  and  $j$ . This suggests that for a sufficiently small scattering of coefficient for every fixed  $k, l$  the quadratic operator will be a contraction. This remark can be expressed as a precise theorem.

The *Lipschitz constant* of an operator  $W : \mathbb{R}^{n+\nu} \rightarrow \mathbb{R}^{n+\nu}$  is

$$L(W) = \sup_{z \neq t} \frac{\|Wz - Wt\|}{\|z - t\|},$$

where  $\|\cdot\|$  is some norm in  $\mathbb{R}^{n+\nu}$ . If this norm can be chosen so that  $L(W) < 1$  then  $W$  will be a strict contraction in this norm with the consequences: unique fixed point, convergence of all trajectories to this point, exponential rate of convergence. Unless otherwise specified, we will use the  $l_1$ -norm in the basis  $e_i^{(f)} = e_i, i = 1, \dots, n$

and  $e_i^{(m)} = e_{n+i}, i = 1, \dots, \nu$  defined as  $\|z\| = \sum_{i=1}^n x_i + \sum_{j=1}^{\nu} y_j$  for  $z = (x, y) = \sum_{i=1}^n x_i e_i^{(f)} + \sum_{j=1}^{\nu} y_j e_j^{(m)}$ .

**Lemma 1.** [2]. Let  $\Delta$  be a convex  $n$ - dimensional compact in  $\mathbb{R}^n$ ,  $F : \Delta \rightarrow \Delta$  be a smooth map. Then (for any norm)  $L(F) \equiv \max_{z \in \Delta} \|d_z F\|$ .

**Lemma 2.** [2]. Let a matrix  $A = (a_{ij})_{i,j=1}^n$  satisfies

$$\sum_{i=1}^n a_{i1} = \sum_{i=1}^n a_{i2} = \dots = \sum_{i=1}^n a_{in}. \quad (11)$$

Then

$$\|A|_{\mathbb{R}_0^n}\| = \frac{1}{2} \max_{j_1 \neq j_2} \sum_{i=1}^n |a_{ij_1} - a_{ij_2}|, \quad (12)$$

where  $A|_{\mathbb{R}_0^n}$  is restriction operator  $A$  on  $\mathbb{R}_0^n$ .

For each  $z \in \mathcal{B}$  we have linear operator  $M_z : \mathcal{B} \rightarrow \mathcal{B}$  defined by  $M_z(t) = zt$ .

**Theorem 1.** The following inequality holds for the Lipschitz's constant

$$\begin{aligned} L(W) \leq & \max_{i_1, i_2, j} \left( \sum_{k=1}^n |p_{i_1 j, k}^{(f)} - p_{i_2 j, k}^{(f)}| + \sum_{l=1}^{\nu} |p_{i_1 j, l}^{(m)} - p_{i_2 j, l}^{(m)}| \right) + \\ & \max_{j_1, j_2, i} \left( \sum_{k=1}^n |p_{i j_1, k}^{(f)} - p_{i j_2, k}^{(f)}| + \sum_{l=1}^{\nu} |p_{i j_1, l}^{(m)} - p_{i j_2, l}^{(m)}| \right). \end{aligned} \quad (13)$$

*Proof.* For the operator  $W$  in  $S$  the derivative is

$$\begin{aligned} d_z W = \frac{1}{2} & \begin{pmatrix} \sum_{j=1}^{\nu} p_{1j,1}^{(f)} y_j & \dots & \sum_{j=1}^{\nu} p_{nj,1}^{(f)} y_j & \sum_{i=1}^n p_{i1,1}^{(f)} x_i & \dots & \sum_{i=1}^n p_{i\nu,1}^{(f)} x_i \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ \sum_{j=1}^{\nu} p_{1j,n}^{(f)} y_j & \dots & \sum_{j=1}^{\nu} p_{nj,n}^{(f)} y_j & \sum_{i=1}^n p_{i1,n}^{(f)} x_i & \dots & \sum_{i=1}^n p_{i\nu,n}^{(f)} x_i \\ \sum_{j=1}^{\nu} p_{1j,1}^{(m)} y_j & \dots & \sum_{j=1}^{\nu} p_{nj,1}^{(m)} y_j & \sum_{i=1}^n p_{i1,1}^{(m)} x_i & \dots & \sum_{i=1}^n p_{i\nu,1}^{(m)} x_i \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ \sum_{j=1}^{\nu} p_{1j,\nu}^{(m)} y_j & \dots & \sum_{j=1}^{\nu} p_{nj,\nu}^{(m)} y_j & \sum_{i=1}^n p_{i1,\nu}^{(m)} x_i & \dots & \sum_{i=1}^n p_{i\nu,\nu}^{(m)} x_i \end{pmatrix} \\ d_z W = 2M_z = 2 & \sum_{k=1}^n x_k M_k^{(f)} + 2 \sum_{l=1}^{\nu} y_l M_l^{(m)}, \end{aligned} \quad (14)$$

where  $M_k^{(f)} = M_{e_k^{(f)}}$  and  $M_l^{(m)} = M_{e_l^{(m)}}$  is the multiplication maps with matrixes  $(p_{ij,k}^{(f)})_{i,k=1}^n$  and respectively  $(p_{ij,l}^{(m)})_{j,l=1}^{\nu}$ .

By Lemma 1 we have  $L(W) = 2 \max_{z \in S} \|M_z\| \leq 2 \max_k \|M_k^{(f)}\| + 2 \max_l \|M_l^{(m)}\|$ .

By Lemma 2,

$$\begin{aligned}\|M_k^{(f)}\| &= \frac{1}{2} \max_{i_1, i_2, j} \left( \sum_{k=1}^n |p_{i_1 j, k}^{(f)} - p_{i_2 j, k}^{(f)}| + \sum_{l=1}^\nu |p_{i_1 j, l}^{(m)} - p_{i_2 j, l}^{(m)}| \right), \\ \|M_l^{(m)}\| &= \frac{1}{2} \max_{j_1, j_2} \left( \sum_{k=1}^n |p_{i j_1, k}^{(f)} - p_{i j_2, k}^{(f)}| + \sum_{l=1}^\nu |p_{i j_1, l}^{(m)} - p_{i j_2, l}^{(m)}| \right).\end{aligned}$$

□

**Corollary 1.** *An evolutionary operator (6) is a strict contraction if*

$$\begin{aligned}\max_{i_1, i_2, j} \left( \sum_{k=1}^n |p_{i_1 j, k}^{(f)} - p_{i_2 j, k}^{(f)}| + \sum_{l=1}^\nu |p_{i_1 j, l}^{(m)} - p_{i_2 j, l}^{(m)}| \right) + \\ \max_{j_1, j_2, i} \left( \sum_{k=1}^n |p_{i j_1, k}^{(f)} - p_{i j_2, k}^{(f)}| + \sum_{l=1}^\nu |p_{i j_1, l}^{(m)} - p_{i j_2, l}^{(m)}| \right) < 1\end{aligned}\quad (15)$$

For evolutionary operators with positive coefficients there is a multiplicative estimate of the distance from the evolutionary operator to the constant one. Let

$$\mu^f \equiv \mu^f(W) = \max_{i_1, i_2, j, k} \frac{p_{i_1 j, k}^{(f)}}{p_{i_2 j, k}^{(f)}}, \quad \mu^m \equiv \mu^m(W) = \max_{i, j_1, j_2, l} \frac{p_{i j_1, l}^{(m)}}{p_{i j_2, l}^{(m)}},$$

and let  $\zeta(W)$  equal to LHS of (15).

**Lemma 3.**

$$\zeta(W) \leq 4 \frac{\mu^f - 1}{\mu^f + 1} + 4 \frac{\mu^m - 1}{\mu^m + 1}. \quad (16)$$

*Proof.* If  $\alpha, \beta > 0$  and  $\mu = \max(\frac{\alpha}{\beta}, \frac{\beta}{\alpha})$  then obviously

$$|\alpha - \beta| = \frac{\mu - 1}{\mu + 1}(\alpha + \beta).$$

Hence

$$|p_{i_1 j, k}^{(f)} - p_{i_2 j, k}^{(f)}| \leq \frac{\mu^f - 1}{\mu^f + 1} (p_{i_1 j, k}^{(f)} + p_{i_2 j, k}^{(f)}), \quad |p_{i_1 j, l}^{(m)} - p_{i_2 j, l}^{(m)}| \leq \frac{\mu^m - 1}{\mu^m + 1} (p_{i_1 j, l}^{(m)} + p_{i_2 j, l}^{(m)}),$$

and respectively

$$|p_{i j_1, k}^{(f)} - p_{i j_2, k}^{(f)}| \leq \frac{\mu^f - 1}{\mu^f + 1} (p_{i j_1, k}^{(f)} + p_{i j_2, k}^{(f)}), \quad |p_{i j_1, l}^{(m)} - p_{i j_2, l}^{(m)}| \leq \frac{\mu^m - 1}{\mu^m + 1} (p_{i j_1, l}^{(m)} + p_{i j_2, l}^{(m)}).$$

It remains to sum these inequalities over  $k$  and respectively over  $l$ , keeping in mind that

$$\sum_{k=1}^n p_{i_1 j, k}^{(f)} = \sum_{k=1}^n p_{i_2 j, k}^{(f)} = \sum_{l=1}^\nu p_{i j_1, l}^{(m)} = \sum_{l=1}^\nu p_{i j_2, l}^{(m)} = 1.$$

□

**Corollary 2.**

$$L(W) \leq 4 \frac{\mu^f - 1}{\mu^f + 1} + 4 \frac{\mu^m - 1}{\mu^m + 1}.$$

**Corollary 3.** *If  $7\mu^f\mu^m - (\mu^f + \mu^m) < 9$  then the evolutionary operator (6) is a strict contraction.*

**Corollary 4.** *Let  $\mu = \max(\mu^f, \mu^m)$ . Then*

$$L(W) \leq 8 \frac{\mu - 1}{\mu + 1}$$

*and if  $\mu < \frac{9}{7}$  then the evolutionary operator (6) is a strict contraction.*

Let us give several examples and check the condition of Corollary 1.

**Example 1.** Consider the operator

$$W : \begin{cases} x'_1 = \frac{3}{7}x_1y_1 + \frac{1}{2}x_1y_2 + \frac{1}{2}x_2y_1 + \frac{4}{7}x_2y_2, \\ x'_2 = \frac{4}{7}x_1y_1 + \frac{1}{2}x_1y_2 + \frac{1}{2}x_2y_1 + \frac{3}{7}x_2y_2, \\ y'_1 = \frac{4}{7}x_1y_1 + \frac{1}{2}x_1y_2 + \frac{1}{2}x_2y_1 + \frac{3}{7}x_2y_2, \\ y'_2 = \frac{3}{7}x_1y_1 + \frac{1}{2}x_1y_2 + \frac{1}{2}x_2y_1 + \frac{4}{7}x_2y_2. \end{cases} \quad (17)$$

The coefficients of the operator (17) as the following

$$\begin{aligned} p_{11,1}^{(f)} &= \frac{3}{7} & p_{12,1}^{(f)} &= \frac{1}{2} & p_{21,1}^{(f)} &= \frac{1}{2} & p_{22,1}^{(f)} &= \frac{4}{7} \\ p_{11,2}^{(f)} &= \frac{4}{7} & p_{12,2}^{(f)} &= \frac{1}{2} & p_{21,2}^{(f)} &= \frac{1}{2} & p_{22,2}^{(f)} &= \frac{3}{7} \\ p_{11,1}^{(m)} &= \frac{4}{7} & p_{12,1}^{(m)} &= \frac{1}{2} & p_{21,1}^{(m)} &= \frac{1}{2} & p_{22,1}^{(m)} &= \frac{3}{7} \\ p_{11,2}^{(m)} &= \frac{3}{7} & p_{12,2}^{(m)} &= \frac{1}{2} & p_{21,2}^{(m)} &= \frac{1}{2} & p_{22,2}^{(m)} &= \frac{4}{7} \end{aligned}$$

It is easy to check that condition (15) satisfied for (17). Indeed,

$$\begin{aligned} & \max_{i_1, i_2, j} \left( \sum_{k=1}^n |p_{i_1 j, k}^{(f)} - p_{i_2 j, k}^{(f)}| + \sum_{l=1}^{\nu} |p_{i_1 j, l}^{(m)} - p_{i_2 j, l}^{(m)}| \right) + \\ & \max_{j_1, j_2, i} \left( \sum_{k=1}^n |p_{i j_1, k}^{(f)} - p_{i j_2, k}^{(f)}| + \sum_{l=1}^{\nu} |p_{i j_1, l}^{(m)} - p_{i j_2, l}^{(m)}| \right) = \frac{4}{7}. \end{aligned}$$

Consequently, this operator is a strict contraction and it has unique fixed point  $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ . Moreover any trajectory of (17) converges to the fixed point.

The following example shows that the condition of Corollary 1 is not satisfied and evolutionary operator has periodic trajectory.

**Example 2.** Consider the operator

$$W : \begin{cases} x'_1 = x_1y_1 \\ x'_2 = x_1y_2 + x_2 \\ y'_1 = x_2y_2 \\ y'_2 = x_1 + x_2y_1 \end{cases} \quad (18)$$

It easy to check that operator (18) does not satisfy the condition of Corollary 1. We rewrite the operator (18) in the form

$$W : \begin{cases} x'_1 = x_1 y_1 \\ y'_1 = (1 - x_1)(1 - y_1) \end{cases} \quad (19)$$

Denote  $x_n = x_1^{(n)}$ ,  $y_n = y_1^{(n)}$  then from (19) we have

$$\begin{cases} x_{n+1} = x_n y_n \\ y_{n+1} = (1 - x_n)(1 - y_n) \end{cases} \quad (20)$$

Since  $0 \leq x_n y_n \leq x_n$  from the first equation of (20) it follows that  $\lim_{n \rightarrow \infty} x_n = x^* = 0$ . Indeed, for  $(x^0, y^0) \in \text{int}(S^1 \times S^1)$  we get from (20)

$$\begin{aligned} \frac{x_{n+2}}{x_{n+1}} &= (1 - x_n) \left( 1 - \frac{x_{n+1}}{x_n} \right), \\ x_{n+2} x_n &= x_{n+1} (1 - x_n) (x_n - x_{n+1}), \\ \lim_{n \rightarrow \infty} x_{n+2} x_n &= \lim_{n \rightarrow \infty} x_{n+1} (1 - x_n) (x_n - x_{n+1}), \\ (x^*)^2 &= 0, \quad x^* = 0. \end{aligned}$$

Now consider the operator

$$W^2 : \begin{cases} x' = xy - x^2 y - xy^2 + x^2 y^2 \\ y' = x + y - xy - x^2 y - xy^2 + x^2 y^2 \end{cases} \quad (21)$$

Clearly, the operator  $W^2$  has fixed points  $(0, y)$ ,  $0 \leq y \leq 1$ . The point  $(0, y)$  is a saddle point.

It is easy to check that the set  $\{(x, y) \in S^1 \times S^1 : x_1 = 0\}$  is an invariant subset for (18). Any point of the invariant subset is periodic point with period two for operator (18). So trajectory of the operator with an initial point form invariant subset does not converge. Thus operator (18) has a trajectory which does non-converge to the fixed point  $(0, 1, \frac{1}{2}, \frac{1}{2})$ .

The following example shows that condition of Corollary 1 is sufficient but is not necessary.

**Example 3.** Consider the operator with coefficients of inheritance

$$\begin{aligned} p_{11,1}^{(f)} &= 0 & p_{12,1}^{(f)} &= 0 & p_{21,1}^{(f)} &= \frac{1}{2} & p_{22,1}^{(f)} &= \frac{1}{2} \\ p_{11,2}^{(f)} &= 1 & p_{12,2}^{(f)} &= 1 & p_{21,2}^{(f)} &= \frac{1}{2} & p_{22,2}^{(f)} &= \frac{1}{2} \\ p_{11,1}^{(m)} &= 0 & p_{12,1}^{(m)} &= \frac{1}{2} & p_{21,1}^{(m)} &= 0 & p_{22,1}^{(m)} &= \frac{1}{2} \\ p_{11,2}^{(m)} &= 1 & p_{12,2}^{(m)} &= \frac{1}{2} & p_{21,2}^{(m)} &= 1 & p_{22,2}^{(m)} &= \frac{1}{2} \end{aligned}$$



i.e. the evolution operator has the form

$$W : \begin{cases} x'_1 = \frac{1}{2}x_2 \\ x'_2 = x_1 + \frac{1}{2}x_2 \\ y'_1 = \frac{1}{2}y_2 \\ y'_2 = y_1 + \frac{1}{2}y_2 \end{cases} \quad (22)$$

It easy to check that operator (22) does not satisfy the condition of Corollary 1.

$$\begin{aligned} & \max_{i_1, i_2, j} \left( \sum_{k=1}^n |p_{i_1 j, k}^{(f)} - p_{i_2 j, k}^{(f)}| + \sum_{l=1}^{\nu} |p_{i_1 j, l}^{(m)} - p_{i_2 j, l}^{(m)}| \right) + \\ & \max_{j_1, j_2, i} \left( \sum_{k=1}^n |p_{i j_1, k}^{(f)} - p_{i j_2, k}^{(f)}| + \sum_{l=1}^{\nu} |p_{i j_1, l}^{(m)} - p_{i j_2, l}^{(m)}| \right) = 2 > 1. \end{aligned}$$

But any trajectory of (22) converges to  $(\frac{1}{3}, \frac{2}{3}, \frac{1}{3}, \frac{2}{3})$ .

Indeed, from (22) we have

$$x_1^{(n+1)} = \frac{1}{2}(1 - x_1^{(n)})$$

We consider following one dimensional dynamical system.

$$f(x) = \frac{1}{2}(1 - x)$$

It has unique fixed point  $x = \frac{1}{3}$  and decreasing on  $[0, 1]$ . Easy to check that  $f'(x) = -\frac{1}{2}$  and  $|f'(\frac{1}{3})| = \frac{1}{2} < 1$  therefore the fixed point  $x = \frac{1}{3}$  is attracting.

We claim that any trajectory of  $f(x)$  converges to the fixed point  $x = \frac{1}{3}$ . Indeed, we have

$$f^n(x) = \sum_{k=1}^n \frac{(-1)^k}{2^k} + (-1)^n \cdot \frac{x}{2^n}$$

and

$$\begin{aligned} \lim_{n \rightarrow \infty} f^{2n}(x) &= \lim_{n \rightarrow \infty} \left( \frac{1}{3} \cdot \frac{2^{2n} - 1}{2^{2n}} + \frac{x}{2^{2n}} \right) = \frac{1}{3}, \\ \lim_{n \rightarrow \infty} f^{2n+1}(x) &= \lim_{n \rightarrow \infty} \left( \frac{1}{3} \cdot \frac{2^{2n} - 1}{2^{2n}} + \frac{1}{2^{2n+1}} - \frac{x}{2^{2n+1}} \right) = \frac{1}{3}. \end{aligned}$$

So for any initial point trajectory of (22) converges to  $(\frac{1}{3}, \frac{2}{3}, \frac{1}{3}, \frac{2}{3})$ .

**Acknowledgment.** The final part of this work was done at the International Islamic University of Malaysia (IIUM) and the second author would like to thank the IIUM for providing financial support and all facilities. The second author also thanks Dr. Mansoor Saburov for useful discussions.

## References

- [1] Bernstein S.N. The solution of a mathematical problem related to the theory of heredity. *Uchn. Zapiski. NI Kaf. Ukr. Otd. Mat.* 1924. no. 1., 83-115 (Russian).
- [2] Lyubich Yu.I. Mathematical structures in population genetics. *Biomathematics*, **22**, Springer-Verlag, 1992.
- [3] Ganikhodzhaev R.N. Quadratic stochastic operators, Lyapunov function and tournaments. *Acad. Sci. Sb.Math.* **76** (1993), no. 2, 489-506.
- [4] Ganikhodzhaev R.N. A chart of fixed points and Lyapunov functions for a class of discrete dynamical systems. *Math. Notes* **56** (1994), no. 5-6, 1125-1131.
- [5] Ganikhodzhaev R.N., Eshmamatova D.B. Quadratic automorphisms of a simplex and the asymptotic behavior of their trajectories. *Vladikavkaz. Mat. Zh.* **8** (2006), no. 2, 12-28 (Russian).
- [6] Ganikhodzhaev N.N. An application of the theory of Gibbs distributions to mathematical genetics. *Dokl. Math.* **61** (2001), 321-323.
- [7] Rozikov U.A., Shamsiddinov N.B. On non-Volterra quadratic stochastic operators generated by a product measure. *Stoch. Anal. Appl.* Vol. **27**, 2 (2009), 353-362.
- [8] Ladra M., Rozikov U.A. Evolution algebra of a bisexual population. *ArXiv*: 0004189 [math.AC]
- [9] Ladra M., Omirov B.A., Rozikov U.A. On dibaric and evolution algebras. *ArXiv*: 1104.2578 [math.RA]
- [10] Rozikov U.A., Zhamilov U.U. On F- quadratic stochastic operators. *Math. Notes* **83** (2008), no. 4, 554-559.
- [11] Zhamilov U.U., Rozikov U.A. On dynamics of stricly non-Volterra quadratic stochastic operators defined on the two dimensional simplex. *Sb. Math.* Vol. **200**. no. 9 (2009), 1339-1351.
- [12] Rozikov U.A., Zhamilov U.U. Volterra quadratic stochastic operators of a bisexual population. *Ukr. Math. Jour.* Vol. **63**. no. 7 (2011), 985-998.(Russian)
- [13] Ganikhodzhaev R.N., Mukhamedov F.M., Rozikov U.A. Quadratic stochastic operators: Results and open problems. *Infinite Dimensional Analysis, Quantum Probability and Related Topics*. Vol. **14**. no. 2 (2011), 279-335.
- [14] Zhamilov U.U., Mukhiddinov R.T. On conditional quadratic stochastic operators. *Uzbek. Math. Zh.* no. 2 (2010), 31-38 (Russian).
- [15] Zhamilov U.U. FM— Volterra quadratic stochastic operators. *Dokl. Acad. Nauk. RUz* no. 3 (2010), 10-14 (Russian).